

PEDIATRIC REFERENCES


Pediatric Vital Signs

Age	Heart Rate	Respiratory Rate	Minimum Systolic BP
Infant (less than 1 year)	100 – 160	30 – 60	greater than 60
Toddler (1 to 2 years)	90 – 150	24 – 40	greater than 70
Preschooler (3 to 5 years)	80 – 140	22 – 34	greater than 75
School-aged child (6 to 10 years)	70 – 120	18 – 30	greater than 80
Adolescent (11 to 18 years)	60 – 100	12 – 16	greater than 90


Pediatric Airway Management Supplies

Weight (kg)	Laryngoscope Blade	ET Tube	ET Tube Length	Stylet	Suction Catheter
Newborn 3-5 kg	0-1 straight	3.0-3.5 uncuffed	10-10.5	6 Fr	6-8 Fr
Infant 6-9 kg	1 straight	3.5 uncuffed	10-10.5	6 Fr	8 Fr
Toddler 10-11 kg	1 straight	4.0 uncuffed	11-12	6 Fr	8-10 Fr
Small Child 12-14 kg	2 straight	4.5 uncuffed	12.5-13.5	6 Fr	10 Fr
Child 15-18 kg	2 straight or curved	5.0 uncuffed	14-15	6 Fr	10 Fr
Child 19-22 kg	2 straight or curved	5.5 uncuffed	15.5-16.5	14 Fr	10 Fr
Large Child 24-30 kg	2-3 straight or curved	6.0 cuffed	17-18	14 Fr	10 Fr
“Adult” greater than or equal to 32 kg	3 straight or curved	6.5 cuffed	18.5-19.5	14 Fr	12 Fr

PEDIATRIC REFERENCES



Wong-Baker FACES Pain Rating Scale



Explain to the person that each face is for a person who feels happy because he has no pain (hurt) or sad because he has some or a lot of pain. Face 0 is very happy because he doesn't hurt at all. Face 1 hurts just a little bit. Face 2 hurts a little more. Face 3 hurts even more. Face 4 hurts a whole lot. Face 5 hurts as much as you can imagine, although you don't have to be crying to feel this bad. Ask the person to choose the face that best describes how he is feeling.

Rating scale is recommended for persons age 3 years and older.