



Class 3 IFT-Paramedic Treatment Protocol 3504

Antivenin

Page 1 of 1

This protocol is used when antivenin has been initiated at the sending facility and must be continued during transport. Antivenin may be used in patients with moderate to severe reaction from a venomous snake bite or a severe reaction from a black widow spider envenomation.

Contraindications: Known allergy. The Crotalidae Polyvalent Antivenin, Antivenin Micrurus Fulvius, and the Black Widow Antivenin are derived from horse/sheep serum, therefore, allergy to horses/sheep and a prior reaction to horse/sheep serum derived products are relative contraindications.

- A. Perform **Class 3 IFTA Protocol 9204**. Perform **Inter-Facility Transport Assessment (IFTA) Procedures Patient Care Protocol 9204** and follow the proper protocol for medical management based on clinical presentation.
- B. Treatment:
1. Maintain infusion rate **ordered by the sending physician**. The IV site should be in the unaffected extremity.
 2. Consider aeromedical transport if available to minimize out of hospital time between the sending and receiving facility.
 3. Monitor vital signs every 15 minutes if asymptomatic and every 5 minutes if symptomatic. Symptoms may include nausea, vomiting, mental status changes, tachycardia, hypotension, and uncontrolled bleeding.

4. Monitor patient for signs and symptoms of allergic or anaphylactic reaction. If symptoms are present, refer to Allergic Reaction/anaphylaxis Protocol 4501 and contact MCP.



5. If symptoms progress or systemic symptoms begin or worsen, **contact Medical Command** for further orders or possible ground CCT intercept.

